arXiv:1611.05227v1 [astro-ph.SR] 16 Nov 2016

Low frequency observations of transient quasi-periodic radio emission from the solar atmosphere

K. Sasikumar Raja[1] and R. Ramesh[1]

sasikumar@iiap.res.in

ABSTRACT

We report low frequency observations of the quasi-periodic, circularly polarized, harmonic type III radio bursts whose associated sunspot active regions were located close to the solar limb. The measured periodicity of the bursts at 80 MHz was ≈ 5.2 s and their average degree of circular polarization (*dcp*) was ≈ 0.12. We calculated the associated magnetic field B : (1) using the empirical relationship between the *dcp* and B for the harmonic type III emission, and (2) from the observed quasi-periodicity of the bursts. Both the methods result in $B \approx 4.2$ G at the location of the 80 MHz plasma level (radial distance $r \approx 1.3$ $R_\odot$) in the active region corona.

Subject headings: Sun: corona — Sun: oscillations — Sun: radio radiation — Sun: magnetic topology

1. Introduction

Type III solar radio bursts are the signatures of accelerated electrons streaming outward through the corona and the interplanetary medium in the aftermath of flares and weak chromospheric brightenings. The electrons stream at speeds $\approx c/3$ along the open field magnetic field lines. The radio emission is widely accepted to be because of the following two-step process: (1) the excitation of high levels of plasma oscillations (Langmuir waves) by the propagating electron streams/beams and (2) subsequent conversion of these Langmuir waves into electromagnetic waves at the fundamental (F) and second harmonic (H) of the local plasma frequency. The H emission results from the coalescence of two Langmuir waves. The type III bursts are best identified on the dynamic spectra obtained using radio spectrographs. They appear as intense emission, drifting from high to low frequencies at

[1]Indian Institute of Astrophysics, II Block, Koramangala, Bangalore - 560 034.

$\approx$ 100 MHz/s. The above frequency drift results from the decrease of electron density and hence the plasma frequency, with distance in the solar atmosphere. The bursts occur over the frequency range from $\approx$ 1 GHz-10 kHz, corresponding to a distance range extending from the low corona to beyond the orbit of the Earth. The bursts occur either isolated or in groups. While the former has a life-time of about few seconds at any given frequency, the groups of bursts last for $\approx$ 5 m (Suzuki & Dulk 1985). Groups of type III bursts are generally associated with flares observed in X-rays and/or Hα. They are a classic signature of the impulsive phase of flares in which the radiation is primarily non-thermal. It is believed that each individual burst is due to radiation associated with electron streams moving outwards through the corona along large scale diverging magnetic field lines connected to a common acceleration/injection site (Mercier 1975; Pick & Ji 1986). The occurence of non-thermal bursts is a clear indication of particle acceleration during an event. Note that the generation of radio emission by non-thermal electron beams is very efficient due to the coherent nature of the emission mechanism. Most of the type III bursts are weakly, circularly polarized with $dcp < 0.15$. However, some bursts, those identified as due to fundamental plasma emission, have $dcp \approx 0.5$ (Suzuki & Sheridan 1978; Dulk & Suzuki 1980). Harmonic emission is observed mainly at low frequencies. Note that the presence of the background magnetic field at the source region of type III radio bursts can give rise to a net *dcp* in the ordinary ('o') mode for the escaping radiation (Melrose & Sy 1972; Melrose, Dulk & Smerd 1978; Melrose, Dulk & Gary 1980; Zlotnik 1981; Willes & Melrose 1997). The polarization observations of the H component of type III radio bursts are considered to be a better diagnostic tool to estimate the solar coronal magnetic field since the polarization of the F component is affected by propagation effects (Dulk & McLean 1978). In this paper we report observations of quasi-periodic, harmonic type III bursts (Wild, Smerd & Weiss 1963; Janssens, White & Broussad 1973; Mangeney & Pick 1989; Zhao, Mangeney & Pick 1991; Aschwanden, Benz & Montello 1994; Ramesh et al. 2003, 2005) and estimate the coronal magnetic field using the empirical relationship between the magnetic field and the *dcp* for the harmonic plasma emission (Melrose, Dulk & Smerd 1978). We verified the results by independantly estimating the magnetic field from the periodicity of the observed type III radio bursts. Interestingly both the methods give consistent results indicating the usefulness of ground based low frequency radio observations to estimate the coronal magnetic field, particularly over the range 1.2 $R_\odot \lesssim r \lesssim 2\ R_\odot$, where measurements at other wavelength bands in the electromagnetic spectrum are presently difficult.

2. Observations

The radio data reported in the present work were obtained at 80 MHz on 2012 September 20, 2013 January 18, and 2013 March 11 with the heliograph (Ramesh et al. 1998, 1999b, 2006), the polarimeter (Ramesh et al. 2008), and the spectrograph (Ebenezer et al. 2007; Kishore et al. 2013) at the Gauribidanur observatory[1], about 100 km north of Bangalore in India (Ramesh 2011a). The co-ordinates of the array are Longitude = $77^{\circ}27^{'}07^{''}$ east, and Latitude = $13^{\circ}36^{'}12^{''}$ North. The heliograph (Gauribidanur RAdioheliograPH, GRAPH) is a T-shaped radio interferometer array which produces two dimensional images of the solar corona with an angular resolution of $\approx 5^{'} \times 7^{'}$ (R.A. $\times$ decl.) at the above frequency. The integration time is $\approx$ 250 ms and the observing bandwidth is $\approx$ 2 MHz. The polarimeter (Gauribidanur Radio Interference Polarimeter, GRIP) is an east-west one-dimensional array operating in the interferometer mode and it responds to the integrated and polarized flux densities from the 'whole' Sun. The half-power width of the response function ('beam') of the GRIP is broad (compared to the Sun) in both right ascension/east-west direction ($\approx 2^{\circ}$ at 80 MHz) and declination/north-south direction ($\approx 90^{\circ}$). This implies that a plot of the GRIP data (i.e. the time profile) for observations in the transit mode is essentially the 'east-west beam' of the array with an amplitude proportional to the strength of the emission from the 'whole' Sun at the observing frequency, weighted by the antenna gain in that direction. The integration time is $\approx$ 250 ms and the observing bandwidth is $\approx$ 2 MHz. The radio source(s) responsible for the circularly polarized emission observed with the GRIP are identified using the two-dimensional radioheliograms obtained with the GRAPH around the same time. Note that linear polarization, if present at the coronal source region, tends to be obliterated at low radio frequencies because of the differential Faraday rotation of the plane of polarization within the observing bandwidth (Grognard & McLean 1973). The spectrograph antenna system (Gauribidanur LOw frequency Solar Spectrograph, GLOSS) is a total power instrument and the half-power width of its antenna response of GLOSS is $\approx 90^{\circ} \times 6^{\circ}$ (R.A. $\times$ decl.) at 80 MHz. The integration time is $\approx$ 100 ms and the observing bandwidth is $\approx$ 300 KHz at each frequency. The width of the response of GLOSS in hour angle is nearly independent of frequency. The Sun is a point source for both the GRIP and the GLOSS. All the above three instruments observe the Sun everyday during the interval 4 - 9 UT. The minimum detectable flux density of the GRAPH, GRIP and GLOSS are $\approx$ 20 Jy, 200 Jy, 3000 Jy, respectively (1 Jy = 10^{-26} $\mathrm{Wm^{-2}Hz^{-1}}$).

Figure 1 shows the temporal evolution of the Stokes I & V radio emission from the solar corona at 80 MHz as observed with the GRIP on 2012 December 20 during the interval 06:36-

[1]http://www.iiap.res.in/centers/radio

06:39 UT, i.e. around the transit of the Sun over Gauribidanur. One can notice the presence of intense quasi-periodic emission in both the Stokes I and V time profiles. These are the characteristic signature of groups of type III solar radio bursts from the Sun (Suzuki & Dulk 1985). We verified the same from the dynamic spectra (85-35 MHz) obtained with GLOSS during the above interval (Figure 2). The peak Stokes I and V flux densities in Figure 1 are $\approx 1.2 \times 10^5$ Jy and 1.6×10^4 Jy, respectively. The corresponding $dcp = \frac{|\text{V flux}|}{\text{I flux}}$ is ≈ 0.11. Note the average flux density of the type III solar radio bursts reported in the literature is typically $\approx 10^7$ Jy (Suzuki & Dulk 1985). This is about two orders of magnitude higher than the peak flux density of the type III burst of 2012 December 20, indicating that the latter is a weak event. The estimated average periodicity of the radio emission in Figure 1 is ≈ 4.5 sec, in both Stokes I & V. The source region of the bursts was identified from the radioheliogram obtained with the GRAPH around the same time (Figures 3 and 4). One can notice the bursts orginated close to the solar limb. They were associated with a SF class Hα flare from AR 11574[2] located at S25W69[3]. A shift in the solar radio source position due to ionospheric effects is expected to be $\lesssim 0.1\ R_\odot$ at 80 MHz in the hour angle range $\pm$ 2 hr (Stewart & McLean 1982). The present observations were carried out close to the transit of the Sun over the local meridian at Gauribidanur. Similarly, the effects of scattering (irregular refraction due to density inhomogeneities in the solar corona) on the observed source position are also considered to be small at 80 MHz compared to lower frequencies (Aubier et al. 1971; Bastian 2004). The positional shift of discrete solar radio sources due to scattering is expected to be $\lesssim 0.2\ R_\odot$ at 80 MHz (Riddle 1974; Robinson 1983; Thejappa et al. 2007). Ray tracing calculations employing realistic coronal electron density models and density fluctuations show that the turning points of the rays that undergo irregular refraction almost coincide with the location of the plasma ('critical') layer in the non-scattering case even at 73.8 MHz (Thejappa & MacDowall 2008). Obviously, the situation should be better at 80 MHz. Also there is no unambiguous evidence that scattering is important in any of the solar radio bursts (McLean & Melrose 1985). Note that high angular resolution observations of the solar corona indicate that discrete radio sources of angular size $\approx 1^{'}$ are present in the solar atmosphere from where low frequency radio radiation originates (Kerdraon 1979; Lang & Willson 1987; Willson et al. 1998; Ramesh et al. 1999b; Ramesh & Sastry 2000; Ramesh & Ebenezer 2001; Mercier et al. 2006; Kathiravan et al. 2011; Ramesh et al. 2012). The projection effects are also expected to be minimal since we have used only the limb events for the present work (see for eg. Figure 4). The details related to the type III radio burst 2012 December 20 described above and the other two events (2013 January 18 and 2013 March 11) are listed

[2]http://www.swpc.noaa.gov/

[3]http://www.lmsal.com/solarsoft/latest_events/

in Table 1.

3. Analysis and Results

3.1. Estimate of B from the estimated dcp of the type III radio bursts

Wild, Sheridan & Nylan (1959) pointed out that both the fundamental (F) and the harmonic (H) type III solar radio burst emission should generally be observable only for events near the center of the solar disk. Elsewhere it should be purely harmonic emission. This is because the F emission is more directive compared to the H emission. According to Caroubalos & Steinberg (1974), the ground based observations of type III radio bursts associated with sunspot regions located $\gtrsim 70^{\circ}$ either to the east or west of the central meridian on the Sun, are primarily H emission. The F component has a limiting directivity of $\pm 65^{\circ}$ (from the central meridian on the Sun) at 80 MHz (Suzuki & Sheridan 1982). A similar result was recently reported by Thejappa, MacDowall & Bergamo (2012) for the very low frequency solar type III radio bursts observed in the interplanetary medium. Note the heliographic longitude of the sunspot regions associated with the type III radio bursts in the present case are all $\gtrsim 70^{\circ}$ (see Table 1). The estimated values of dcp for all the three events in the present case are close to the average dcp (≈ 0.11) reported for the circularly polarized H component of the type III bursts (see Table 1). For the F component, the reported average $dcp \approx 0.35$ (Dulk & Suzuki 1980). The above arguments on the directivity and the dcp indicate that the type III bursts observed in the present case are due to H emission. Under such circumstances the magnetic field (B) near the source region of the bursts can be calculated using the following relationship (Suzuki & Sheridan 1978; Dulk & Suzuki 1980; Mercier 1990; Reiner et al. 2007; Melrose, Dulk & Smerd 1978; Zlotnik 1981; Ramesh et al. 2010b):

$$B = \frac{f_p \times dcp}{2.8\ a(\theta, \theta_0)} \tag{1}$$

where f_p is the plasma frequency in MHz and B is in Gauss. $a(\theta, \theta_0)$ is a slowly varying function which depends on the viewing angle θ between the magnetic field component and the line-of-sight, and the angular distribution θ_0 of the Langmuir waves. In the present case, $\theta \approx 70^{\circ} - 90^{\circ}$ (see Table 1). We assumed $f_p = 40$ MHz since the observed emission at 80 MHz is most likely the H component. The Langmuir waves are considered to be confined to a small range of angles, i.e. a cone of opening angle $\theta_0 \approx 10^{\circ} - 30^{\circ}$, to the magnetic field direction for harmonic type III emission in the 'o' mode (Melrose, Dulk & Smerd 1978; Dulk & Suzuki 1980; Willes & Melrose 1997; Benz 2002). Assuming the average value, i.e. $\theta_0 \approx 20^{\circ}$, we find that $a(\theta, \theta_0) \approx 0.4$ in the aforementioned range of θ (Melrose, Dulk & Gary

1980; Gary 1982; Suzuki & Dulk 1985). The estimated magnetic field values using equation (1) for the type III events reported in the present work are listed in Table 1. Figure 5 shows the B values corresponding to the individual bursts in the quasi-periodic type III burst emission observed on 20 September 2012 (Figure 1). They remain approximately constant, within the error limits. We found that the B values corresponding to the bursts observed on 2013 January 18 and 2013 March 11 also exhibit a similar trend.

3.2. Estimate of B from the observed quasi-periodicity of the type III radio bursts

The term quasi-periodicity or pulsations in solar radio physics refers to the quasi-periodic amplitude variations in the time profile of the observed radio flux at any particular observing frequency. The associated physical mechanisms have been classified into three categories as on date: (1) modulation of the emission by coronal loop oscillations, (2) intrinsic oscillations of the emission created by oscillatory wave-wave interactions and wave-particle interactions, and (3) modulation of electron acceleration/injection process responsible for the emission (Aschwanden 1987; Nindos & Aurass 2007). Out of this, category (2) can be ruled out in the present case since they apply primarily to fundamental 'o' mode and harmonic extraordinary mode ('e' mode) whereas the type III bursts reported in the present work correspond to harmonic 'o' mode as mentioned in Section 3.1 (Aschwanden & Benz 1988). In view of this, we have carried out the calculations for categories (1) and (3) in the reminder of this section. Laboratory experiments and numerical simulations have shown that the release of magnetic energy by reconnection must be considered as a highly time-dependent process (see for eg. Pick & van den Oord (1990)). According to Kliem, Karlicky & Benz (2000), the temporal variations in the radio burst flux are caused by modulations of the particle acceleration in a highly dynamic reconnection process. A similar model, i.e. quasi-periodic reconnection and particle injection was reported by Zlotnik et al. (2003) also. The authors had noted that even a very small ($\approx 2\%$) quasi-periodic modulation of the magnetic field is sufficient for periodic electron acceleration. Both these models belong to the category (3) mentioned above and were successful in explaining the associated observations. The modulation is likely to be communicated on a magnetohydrodynamic (MHD) time scale in the acceleration region (Tajima et al. 1987; Aschwanden, Benz & Montello 1994; Kliem, Karlicky & Benz 2000; Asai et al. 2001). In such a case, the corresponding Alfvén speed (v_A) can be estimated as:

$$v_A \approx l/p \tag{2}$$

where p is the period of the quasi-periodic emission in seconds (s) and $l \approx 10000$ km is the typical dimension of the region over which the type III radio burst producing elec-

trons are injected (Lantos, Pick & Kundu 1984; Aschwanden 2002). It is possible that the individual bursts that are temporally separated in a type III group are also spatially fragmented within the same acceleration region (Pick & van den Oord 1990; Vlahos & Raoult 1995; Isliker et al. 1998). Considering category (1) where the observed quasi-periodicities are caused by the MHD oscillations of the associated coronal loop(s), the corresponding relationship for the Alfvén speed is (Roberts, Edwin & Benz 1984),

$$v_A = 2.6a/p \tag{3}$$

where $a \approx 3500$ km is the width of the coronal loop. According to Aschwanden (1987), the quasi-periodic emission observed in the upper part of the corona from where the meter-decameter wave radio emission originates, paticularly those due to plasma processes as in the present case, are best described by the above model. Asai et al. (2001) showed that the quasi-periodic pulsation observed by them in the microwave range are due to the modulation of the acceleration/injection rate of the non thermal electrons by the oscillations in the associated coronal loop(s). Once v_A is known in either category (1) or (3), the associated magnetic field (B) can be calculated from the definition of the former, i.e.:

$$v_A = 2.05 \times 10^6 \; B \; N_e^{-1/2} \tag{4}$$

N_e is the electron density in units of cm^{-3} and can be estimated using the relationship $f_p = 9 \times 10^{-3} N_e^{1/2}$. Note v_A in equation (3) is in units of km/s. The v_A and B values estimated using equations (2) and (4) for the type III events reported in the present work are listed in Table 1. Equations (3) and (4) result in nearly the same values for v_A and B.

4. Summary

We have reported estimates of the coronal magnetic field using low frequency (80 MHz) radio observations of quasi-periodic harmonic type III burst emission associated with sunspot regions close to the solar limb. The data were obtained with the heliograph, the polarimeter and the spectrograph at the Gauribidanur observatory. Though several type III bursts are observed everyday with the aforementioned suite of instruments, we selected only those events with $dcp \lesssim 0.15$ and whose associated source region is close to the limb of the Sun. The above are the criteria to identify a harmonic type III burst as mentioned earlier. Also we limited ourselves to data obtained close to the transit of the Sun over the local meridian at Gauribidanur in order to minimize errors in the source position due to propagation effects. We adopted two independant approaches to estimate the magnetic field: (1) based on the relationship between the dcp and the B for harmonic type III radio burst emission, and (2) using the quasi-periodicity in the observed type III radio burst emission. Both the methods

give the same result, i.e. the average $B \approx 4.2$ G. We would like to note here that the often referred empirical relationship for the coronal magnetic field (Dulk & McLean 1978) predicts $B \approx 3$ G at 80 MHz. We had assumed the 80 MHz plasma level to be located at radial distance $r \approx 1.3$ $R_\odot$ in the solar atmosphere for the above calculation (Ramesh et al. 2011b). Note Lin, Kuhn & Coulter (2004) had measured $B \approx 4$ G at $r \approx 1.1$ $R_\odot$ using Zeeman splitting observations of the Fe XIII λ1075 nm coronal emission line from an active region close to the solar limb. Assuming that $B \propto r^{-1.5}$ (Dulk & McLean 1978), we find that the above measurement indicates $B \approx 3.1$ G $r \approx 1.3$ $R_\odot$. The present estimates are also in reasonable agreement with that of Ramesh et al. (2010a) who had reported $B \approx 5 \pm 1$ G at 77 MHz for radio sources associated with the coronal streamers at $r \approx 1.5$ $R_\odot$. Note the type III radio bursts are considered to be closely associated with the coronal streamers (Kundu et al. 1983). We would like to add here that the estimated v_A in the present case (see Table 1) are consistent with the corresponding values reported by Gopalswamy et al. (2001); Vršnak et al. (2002) for the active region solar corona.

It is a pleasure to thank the staff of the Gauribidanur observatory for their help in observations, maintenance of the antenna and receiver systems there. We thank the referee for his/her kind suggestions and comments using which we were able to bring out the results more clearly.

REFERENCES

Asai, A., Shimojo, M., Isobe, H., Morimoto, T., Yokoyama, T., Shibasaki, K. & Nakajima, H. 2001, ApJ, 562, L103

Aschwanden, M. J. 1987, Sol. Phys., 111, 113

Aschwanden, M. J. & Benz, A. O. 1988, ApJ, 332, 466

Aschwanden, M. J., Benz, A. O. & Montello, M. L. 1994, ApJ, 431, 432

Aschwanden, M. J. 2002, Space Sci. Rev., 101, 178

Aubier, M., Leblanc, Y. & Boischot, A. 1971, A&A, 12, 435

Bastian, T. S. 2004, Planet. Sp. Sci., 52, 1381

Benz, A. O. 2002, Plasma Astrophysics. Kinetic Processes in Solar and Stellar Coronae (Dordrecht: Kluwer), Astrophys. Sp. Sci. Lib., 279

Caroubalos, C. & Steinberg, J. -L. 1974, in IAU Symp. 57, Coronal Disturbances, ed. G. A. Newkirk (Dordrecht: Reidel), 239

Dulk, G. A. & McLean, D. J. 1978, Sol. Phys., 57, 279

Dulk, G. A. & Suzuki, S. 1980, A&A, 88, 203

Ebenezer, E., Subramanian, K. R., Ramesh, R., Sundara Rajan, M. S. & Kathiravan, C. 2007, Bull. Astron. Soc. India, 35, 111

Gary, D. E. 1982, Radio Emission from Solar and Stellar Coronae, Ph.D Thesis, University of Colorado at Boulder, (Ann Arbor: University Microfilms International), 87

Gopalswamy, N., Lara, A., Kaiser, M. L. & Bougeret, J. -L. 2001, J. Geophys. Res., 106(A11), 25261

Grognard, R. J.-M. & McLean, D. J. 1973, Sol. Phys., 29, 149

Isliker, H., Vlahos, L., Benz, A. O. & Raoult, A. 1998, A&A, 336, 371

Janssens, T. J., White, K. P. & Broussad, R. M. III 1973, Sol. Phys., 31, 207

Kathiravan, C., Ramesh, R., Indrajit V. Barve & Rajalingam, M. 2011, ApJ, 730, 91

Kerdraon, A. 1979, A&A, 71, 266

Kishore, P., Kathiravan, C., Ramesh, R., Indrajit V. Barve & Rajalingam, M. 2013, (in preparation)

Kliem, B. Karlićky, M., & Benz, A. O. 2003, A&A, 360, 715

Kundu, M. R., Gergely, T. E., Turner, P. J. & Howard, R. A. 1983, ApJ, 269, L67

Lang, K. R., & Willson, R. F. 1987, ApJ, 319, 514

Lantos, P., Pick, M. & Kundu, M.R. 1984, ApJ, 283, L71

Lin, H., Kuhn, J. R. & Coulter, R. 2004, ApJ, 613, L177

Mangeney, A. & Pick, M. 1989, A&A, 224, 242

McLean, D. J. & Melrose, D. B. 1985, in Solar Radio Physics, ed. D.J. McLean & N.R. Labrum, (Cambridge: Cambridge University Press), 237

Melrose, D. B., & Sy, W. N. 1972, Aust. J. Phys., 25, 387

Melrose, D. B., Dulk, G. A. & Smerd, S. F. 1978, A&A, 66, 315

Melrose, D. B., Dulk, G. A. & Gary, D. E. 1980, Proc. Astron. Soc. Aust., 4, 50

Mercier, C. 1975, Sol. Phys., 45, 169

Mercier, C. 1990, Sol. Phys., 130, 119

Mercier, C., et al. 2006, A&A, 447, 1189

Nindos, A. & Aurass, H. 2007, in The High Energy Solar Corona: Waves, Eruptions, Particles, ed. K. -L. Klein & A. L. Mackinnon, Lecture Notes in Physics (Berlin: Springer-Verlag), 725, 251

Pick, M. & Ji, S. C. 1986, Sol. Phys., 107, 159

Pick, M. & van den Oord, G. H. J. 1990, Sol. Phys., 130, 83

Ramesh, R., Subramanian, K. R., Sundara Rajan, M. S. & Sastry, Ch. V. 1998, Sol. Phys., 181, 439

Ramesh, R., Subramanian, K. R. & Sastry, Ch. V. 1999a, Sol. Phys., 185, 77

Ramesh, R., Subramanian, K. R. & Sastry, Ch. V. 1999, A&AS, 139, 179

Ramesh, R. & Sastry, Ch. V. 2000, A&A, 358, 749

Ramesh, R. & Ebenezer, E. 2001, ApJ, 558, L141

Ramesh, R., Kathiravan, C., Satya Narayanan, A. & Ebenezer, E. 2003, A&A, 400, 753

Ramesh, R., Satya Narayanan, A., Kathiravan, C., Sastry, Ch. V. & Udaya Shankar, N. 2005, A&A, 431, 353

Ramesh, R., Sundara Rajan, M. S. & Sastry, Ch. V. 2006, Exp. Astron., 21, 31

Ramesh, R., Kathiravan, C., Sundara Rajan, M. S., Indrajit V. Barve & Sastry, Ch. V. 2008, Sol. Phys., 253, 319

Ramesh, R. Kathiravan, C. & Sastry, Ch. V. 2010a, ApJ, 711, 1029

Ramesh, R., Kathiravan, C., Indrajit V. Barve, Beeharry, G. K. & Rajasekara G. N. 2010b, ApJ, 719, L41

Ramesh, R. 2011a, in Proc. Astron. Soc. India Conf. Sec. 2, 1st Asia-Pacific Solar Physics Meeting, ed. A. R. Choudhuri & D. Banerjee (Banglore: ASI), 55

Ramesh, R., Kathiravan, C. & Satya Narayanan, A. 2011b, ApJ, 734, 39

Ramesh, R., Kathiravan, C., Indrajit V. Barve & Rajalingam, M. 2012, ApJ, 744, 165

Reiner, M. J., Fainberg, J., Kaiser, M. L. & Bougeret, J. -L. 2007, Sol. Phys., 241, 351

Riddle, A. C. 1974, Sol. Phys., 35, 153

Robinson, R. D. 1983, Proc. Astron. Soc. Aust., 5, 208

Roberts, B., Edwin, P. M. & Benz, A. O. 1984, ApJ, 279, 857

Stewart, R. T. & McLean, D. J. 1982, Proc. Astron. Soc. Aust., 4, 386

Suzuki, S. & Sheridan, K. V. 1978, Quantum Electron., 20, 989

Suzuki, S. & Sheridan, K. V. 1982, Proc. Astron. Soc. Aust., 4, 382

Suzuki, S. & Dulk, G. A. 1985, in Solar Radio Physics, ed. D.J. McLean & N.R. Labrum, (Cambridge: Cambridge University Press), 289

Tajima, T., Sakai, J., Nakajima, H., Kosugi, T., Brunel, F. & Kundu, M. R. 1987, ApJ, 321, 1031

Thejappa, G., MacDowall, R. J. & Kaiser, M. L. 2007, ApJ, 671, 894

Thejappa, G., & MacDowall, R. J. 2008, ApJ, 676, 1338

Thejappa, G., MacDowall, R. J. & Bergamo, M. 2012, ApJ, 745, 187

Vlahos, L. & Raoult, A. 1995, A&A, 296, 844

Vršnak, B., Magdalenić, J., Aurass, H. & Mann, G. 2002, A&A, 396, 673

Wild, J. P., Sheridan, K. V. & Nylan, A. A. 1959, Aust. J. Phys., 12, 369

Wild, J. P., Smerd, S. F. & Weiss, A. A. 1963, ARA&A, 1, 291

Willes, A. J. & Melrose, D. B. 1996, Sol. Phys., 171, 393

Willson, R. F., Redfield, S. L., Lang, K. R., Thompson, B. J., & St. Cyr., O. C. 1998, ApJ, 504, L117

Zhao, R.-Y., Mangeney, A. & Pick, M. 1991, A&A, 241, 183

Zlotnik, E. Ia. 1981, A&A, 101, 250

Zlotnik, E. Ya., Zaitsev, V. V., Aurass, H., Mann, G. & Hofmann, A. 2003, A&A, 410, 1011

This preprint was prepared with the AAS LaTeX macros v5.2.



Fig. 1.— GRIP observations of group of type III solar radio bursts on 2012 September 20 at 80 MHz in Stokes I (upper panel) and Stokes V (lower panel). The arrow marks indicate the individual bursts.



Fig. 2.— GLOSS dynamic spectra (85-35 MHz) of the group of type III solar radio bursts in Figure 1.



Fig. 3.— GRAPH radioheliogram obtained on 2012 September 20 around 06:30 UT, prior to the quasi-periodic type III burst emission in Figure 1. The open circle at the center represents the solar limb. The size of the GRAPH beam at 80 MHz is shown near the lower right. The intense discrete source close to the west limb is the source region of the type III bursts in Figures 1 and 2.



Fig. 4.— Same as Figure 3 but obtained during the peak phase of the burst in Figure 1 at ≈ 06:38 UT.



Fig. 5.— B values corresponding to the individual type III bursts in the quasi-periodic emission in Figure 1, based on the relationship between B and dcp.

Table 1. Paramaters related to the group of type III radio bursts observed with the Gauribidanur facilities

S.No .	Date	Time (UT)	Period p (s)	Alfvén speed v_A (km/s)	Sunspot heliographic co-ordinates	Viewing angle θ (deg)	dcp	Magnetic field B (G)	
								Harmonic emission[a]	Quasi-periodic emission[b]
1	20 Sep 2012	06:35-06:38	4.5	2222	S25W69	71	0.13	4.7 ± 1	4.8 ± 0.3
2	18 Jan 2013	06:51-06:57	6.4	1562	N18W88	88	0.08	2.9 ± 1	3.4 ± 0.3
3	11 Mar 2013	07:00-07:03	4.8	2083	N10E88	88	0.14	5.0 ± 1	4.5 ± 0.3

[a]See Section 3.1 for details.

[b]See Section 3.2 for details.